

TRANSLATOR'S DECLARATION

I, Janet Hope, BSc(Hons.), MIL., MITI., translator to Messrs. Taylor and Meyer of 20 Kingsmead Road, London, SW2 3JD, Great Britain, verify that I know well both the German and the English language, that I have prepared the attached English translation of 39 pages of a German Patent application in the German language with the title:

Verfahren zur fermentativen Herstellung von D-Pantothensäure unter Verwendung coryneformer Bakterien

identified by the code number 000439 BT at the upper left of each page and that the attached English translation of this document is a true and correct translation of the document attached thereto to the best of my knowledge and belief.

I further declare that all statements made of my own knowledge are true and that all statements made on information and belief are believed to be true, and further that these statements are made with the knowledge that wilful false statements and the like are punishable by fine or imprisonment, or both, under 18 USC 1001, and that such false statements may jeopardize the validity of this document.

Signed:

Dated: 15th March 2004

FEDERAL REPUBLIC OF GERMANY

Certificate of Priority for Filing of a Patent Application

Filing number: 100 48 604.5

Filing date: 30th September 2000

Applicant/Proprietor: Degussa AG, Düsseldorf/Germany

First applicant: Degussa-Hüls Aktiengesellschaft,
Frankfurt am Main/Germany

Title: Process for the fermentative preparation of D-
pantothenic acid using coryneform bacteria

IPC: C 12 N, C 07 H, C 12 P

The attached papers are a true and accurate reproduction of the original documents for this patent application.

Munich, 7th August 2001

**On behalf of the President of the German
Patent and Trade Mark Office**

(signature)

Dzierzon

Process for the fermentative preparation of D-pantothenic acid using coryneform bacteria

The invention provides a process for the fermentative preparation of D-pantothenic acid using coryneform bacteria
5 in which the poxB gene is attenuated.

Prior art

Pantothenic acid is a vitamin of commercial importance which is used in human medicine and in the pharmaceuticals industry, in the foodstuffs industry and very particularly
10 in animal nutrition.

Pantothenic acid can be prepared by chemical synthesis, or biotechnologically by fermentation of suitable microorganisms in suitable nutrient solutions. In the chemical synthesis, DL-pantolactone is an important
15 intermediate stage. It is prepared in a multi-stage process from formaldehyde, isobutylaldehyde and cyanide. In further process steps, the racemic mixture is separated, D-pantolactone is subjected to a condensation reaction with β -alanine, and the desired D-pantothenic acid is obtained
20 in this way.

The advantage of the fermentative preparation by microorganisms lies in the direct formation of the desired stereoisomeric D-form, which is free from L-pantothenic acid.

25 Various types of bacteria, such as e.g. Escherichia coli (E. coli), Arthrobacter ureafaciens, Corynebacterium erythrogenes, Brevibacterium ammoniagenes, and also yeasts, such as e.g. Debaromyces castellii, can produce D-pantothenic acid in a nutrient solution which comprises
30 glucose, DL-pantoic acid and β -alanine, as shown in EP-A 0 493 060. EP-A 0 493 060 furthermore shows that in the case of E. coli, the formation of D-pantothenic acid is improved by amplification of pantothenic acid biosynthesis

genes from E. coli which are contained on the plasmids pFV3 and pFV5 in a nutrient solution comprising glucose, DL-pantoic acid and β -alanine.

EP-A 0 590 857 and US Patent 5,518,906 describe mutants
5 derived from E. coli strain IF03547, such as FV5714, FV525, FV814, FV521, FV221, FV6051 and FV5069, which carry resistances to various antimetabolites, such as salicylic acid, α -ketobutyric acid, β -hydroxyaspartic acid, O-methylthreonine and α -ketoisovaleric acid. They produce
10 pantoic acid in a nutrient solution comprising glucose, and D-pantothenic acid in a nutrient solution comprising glucose and β -alanine. It is furthermore shown in EP-A 0 590 857 and US Patent 5,518,906 that after amplification of the pantothenic acid biosynthesis genes contained on the
15 plasmid pFV31, in the abovementioned strains the production of D-pantoic acid in nutrient solutions comprising glucose and the production of D-pantothenic acid in a nutrient solution comprising glucose and β -alanine is improved.

Processes for the preparation of D-pantothenic acid with
20 the aid of Corynebacterium glutamicum (C. glutamicum) are known only in some instances in the literature. Sahm and Eggeling (Applied and Environmental Microbiology 65(5), 1973-1979 (1999)) thus report on the influence of over-expression of the panB and panC genes and Dusch et al.
25 (Applied and Environmental Microbiology 65(4), 1530-1539 (1999)) report on the influence of the panD gene on the formation of D-pantothenic acid.

Object of the invention

The inventors had the object of providing new principles
30 for improved processes for the fermentative preparation of pantothenic acid with coryneform bacteria.

Description of the invention

When D-pantothenic acid or pantothenic acid or pantothenate are mentioned in the following text, this means not only the free acids but also the salts of D-pantothenic acid, such as e.g. the calcium, sodium, ammonium or potassium salt.

The invention provides a process for the fermentative preparation of D-pantothenic acid using coryneform bacteria in which the nucleotide sequence which codes for the enzyme pyruvate oxidase (EC 1.2.2.2) (poxB gene) is attenuated.

This invention also provides a process for the fermentative preparation of D-pantothenic acid, in which the following steps are carried out:

- a) fermentation of D-pantothenic acid-producing coryneform bacteria in which at least the nucleotide sequence which codes for pyruvate oxidase (EC 1.2.2.2) (poxB) is attenuated, in particular eliminated;
- b) concentration of the D-pantothenic acid in the medium or in the cells of the bacteria; and
- c) isolation of the D-pantothenic acid produced.

The strains employed optionally already produce D-pantothenic acid before attenuation of the poxB gene.

Preferred embodiments are to be found in the claims.

The term "attenuation" in this connection describes the reduction or elimination of the intracellular activity of one or more enzymes (proteins) in a microorganism which are coded by the corresponding DNA, for example by using a weak promoter or using a gene or allele which codes for a corresponding enzyme (protein) with a low activity or inactivates the corresponding gene or enzyme (protein), and optionally combining these measures.

The microorganisms which the present invention provides can produce D-pantothenic acid from glucose, sucrose, lactose, fructose, maltose, molasses, starch, cellulose or from glycerol and ethanol. They are representatives of

5 coryneform bacteria, in particular of the genus *Corynebacterium*. Of the genus *Corynebacterium*, there may be mentioned in particular the species *Corynebacterium glutamicum*, which is known among experts for its ability to produce L-amino acids.

10 Suitable strains of the genus *Corynebacterium*, in particular of the species *Corynebacterium glutamicum*, are, for example, the known wild-type strains

Corynebacterium glutamicum ATCC13032

Corynebacterium acetoglutamicum ATCC15806

15 *Corynebacterium acetoacidophilum* ATCC13870

Corynebacterium melassecola ATCC17965

Corynebacterium thermoaminogenes FERM BP-1539

Brevibacterium flavum ATCC14067

Brevibacterium lactofermentum ATCC13869 and

20 *Brevibacterium divaricatum* ATCC14020

and D-pantothenic acid-producing mutants prepared therefrom, such as, for example

Corynebacterium glutamicum ATCC13032 Δ ilvA/pEC7panBC

Corynebacterium glutamicum ATCC13032/pND-D2

25 It has been found that coryneform bacteria produce pantothenic acid in an improved manner after attenuation of the *poxB* gene, which codes for pyruvate oxidase (EC 1.2.2.2).

"Isolated" means separated out of its natural environment.

30 "Polynucleotide" in general relates to polyribonucleotides and polydeoxyribonucleotides, it being possible for these to be non-modified RNA or DNA or modified RNA or DNA.

The nucleotide sequence of the poxB gene is shown in SEQ ID No. 1 and the resulting amino acid sequence of the enzyme protein is shown in SEQ ID No. 2.

5 The poxB gene described in SEQ ID No. 1 can be used according to the invention. Alleles of the poxB gene which result from the degeneracy of the genetic code or due to "sense mutations" of neutral function can furthermore be used.

10 A new nucleotide sequence, shown in SEQ ID No. 6, which lies upstream of the nucleotide sequence of the poxB gene region shown in SEQ ID No. 1 has been found. A new nucleotide sequence, shown in SEQ ID No. 7, which lies downstream of the nucleotide sequence of the poxB gene region shown in SEQ ID No. 1 has furthermore been found.
15 The sequence of the poxB gene region shown in SEQ ID No. 4 has been obtained in this manner.

It has been found that these polynucleotides shown in SEQ ID No. 6 and 7 are useful in the production of mutants with an attenuated, in particular eliminated, poxB gene.

20 It has also been found that coryneform bacteria produce pantothenic acid in an improved manner after attenuation of the poxB gene.

To achieve an attenuation, either the expression of the poxB gene or the catalytic properties of the enzyme protein
25 can be reduced or eliminated. The two measures can optionally be combined.

The decrease in gene expression can take place by suitable culturing or by genetic modification ("mutation") of the signal structures of gene expression. Signal structures of
30 gene expression are, for example, repressor genes, activator genes, operators, promoters, attenuators, ribosome binding sites, the start codon and terminators. The expert can find information on this e.g. in the patent

application WO 96/15246, in Boyd and Murphy (Journal of Bacteriology 170: 5949 (1988)), in Voskuil and Chambliss (Nucleic Acids Research 26: 3548 (1998), in Jensen and Hammer (Biotechnology and Bioengineering 58: 191 (1998)),
5 in Patek et al. (Microbiology 142: 1297 (1996)) and in known textbooks of genetics and molecular biology, such as e.g. the textbook by Knippers ("Molekulare Genetik", 6th edition, Georg Thieme Verlag, Stuttgart, Germany, 1995) or that by Winnacker ("Gene und Klone", VCH
10 Verlagsgesellschaft, Weinheim, Germany, 1990).

Mutations which lead to a change or reduction in the catalytic properties of enzyme proteins are known from the prior art. Examples which may be mentioned are the works of Qiu and Goodman (Journal of Biological Chemistry 272: 8611-
15 8617 (1997)), Sugimoto et al. (Bioscience Biotechnology and Biochemistry 61: 1760-1762 (1997)) and Möckel ("Die Threonindehydratase aus Corynebacterium glutamicum: Aufhebung der allosterischen Regulation und Struktur des Enzyms", Reports from the Jülich Research Centre, Jül-2906,
20 ISSN09442952, Jülich, Germany, 1994). Summarizing descriptions can be found in known textbooks of genetics and molecular biology, such as e.g. that by Hagemann ("Allgemeine Genetik", Gustav Fischer Verlag, Stuttgart, 1986).

25 Possible mutations are transitions, transversions, insertions and deletions. Depending on the effect of the amino acid exchange on the enzyme activity, "missense mutations" or "nonsense mutations" are referred to. Insertions or deletions of at least one base pair (bp) in a
30 gene lead to "frame shift mutations", which lead to incorrect amino acids being incorporated or translation being interrupted prematurely. Deletions of several codons typically lead to a complete loss of the enzyme activity. Instructions on generation of such mutations are prior art
35 and can be found in known textbooks of genetics and

molecular biology, such as e.g. the textbook by Knippers ("Molekulare Genetik", 6th edition, Georg Thieme Verlag, Stuttgart, Germany, 1995), that by Winnacker ("Gene und Klone", VCH Verlagsgesellschaft, Weinheim, Germany, 1990) or that by Hagemann ("Allgemeine Genetik", Gustav Fischer Verlag, Stuttgart, 1986).

An example of a plasmid with the aid of which an insertion mutagenesis of the *poxB* gen can be carried out is pCR2.1poxBint (figure 1).

10 Plasmid pCR2.1poxBint comprises the plasmid pCR2.1-TOPO described by Mead et al. (Bio/Technology 9:657-663 (1991)), into which an internal fragment of the *poxB* gene, shown in SEQ-ID No. 3 has been incorporated. After transformation and homologous recombination in the chromosomal *poxB* gene
15 (insertion), this plasmid leads to a total loss of the enzyme function.

Another example of a mutated *poxB* gene is the Δ poxB allele contained in the plasmid pCRB1-poxBdel (figure 2). The Δ poxB allele contains only the 5' and the 3' flank of the
20 *poxB* gene. The 1737 bp long section of the coding region is missing (deletion). The nucleotide sequence of the Δ poxB allele and of the 5' and 3' flank is shown in SEQ ID No. 12. This Δ poxB allele can be incorporated into coryneform bacteria by integration mutagenesis. The abovementioned
25 plasmid pCRB1-poxBdel is used for this, or the Δ poxB allele is transferred to the plasmid pK18mobsacB and the plasmid of the type pK18mobsacBpoxBdel thereby formed is used. After transfer by conjugation or transformation and homologous recombination by means of a first "cross-over"
30 event which effects integration and a second "cross-over" event which effects excision in the *poxB* gene, the incorporation of the Δ poxB allele is achieved and a total loss of the enzyme function in the particular strain is achieved. The invention provides the Δ poxB allele
35 characterized by SEQ ID No. 12.

Further instructions and explanations on insertion mutagenesis or integration mutagenesis and gene replacement are to be found, for example, in Schwarzer and Pühler (Bio/Technology 9,84-87 (1991)), Peters-Wendisch et al. (Microbiology 144, 915-927 (1998)) or Fitzpatrick et al. (Applied Microbiology and Biotechnology 42, 575-580 (1994)).

It may furthermore be advantageous for the production of pantothenic acid, in addition to the attenuation of the gene which codes for pyruvate oxidase, for one or more of the genes chosen from the group consisting of

- the panB gene which codes for ketopantoate hydroxymethyl transferase (Sahm et al., Applied and Environmental Microbiology, 65, 1973-1979 (1999)),
- 15 • the panC gene which codes for pantothenate synthetase (Sahm et al., Applied and Environmental Microbiology, 65, 1973-1979 (1999)),
- the ilvC gene which codes for acetohydroxy-acid isomeroreductase (EMBL gene library: Accession No. L09232), and
- 20 • the ilvD gene which codes for dihydroxy-acid dehydratase (EP-A-1006189);

to be enhanced, in particular over-expressed.

The term "enhancement" in this connection describes the increase in the intracellular activity of one or more enzymes in a microorganism which are coded by the corresponding DNA, for example by increasing the number of copies of the gene or genes, using a potent promoter or using a gene which codes for a corresponding enzyme having a high activity, and optionally combining these measures.

It may furthermore be advantageous for the production of pantothenic acid, in addition to the attenuation of the gene which codes for pyruvate oxidase, for the pck gene which codes for phosphoenol pyruvate carboxykinase (PEP carboxykinase) (DE: 19950409.1, DSM 13047) to be attenuated.

Finally, in addition to attenuation of pyruvate oxidase, it may be advantageous for the production of pantothenic acid to eliminate undesirable side reactions (Nakayama: "Breeding of Amino Acid Producing Micro-organisms", in: Overproduction of Microbial Products, Krumphanzl, Sikyta, Vanek (eds.), Academic Press, London, UK, 1982) which reduce the production of pantothenic acid.

The microorganisms prepared according to the invention can be cultured continuously or discontinuously in the batch process (batch culture) or in the fed batch (feed process) or repeated fed batch process (repetitive feed process) for the purpose of pantothenic acid production. A summary of known culture methods are described in the textbook by Chmiel (Bioprozesstechnik 1. Einführung in die Bioverfahrenstechnik (Gustav Fischer Verlag, Stuttgart, 1991)) or in the textbook by Storhas (Bioreaktoren und periphere Einrichtungen (Vieweg Verlag, Braunschweig/Wiesbaden, 1994)).

The culture medium to be used must meet the requirements of the particular microorganisms in a suitable manner. Descriptions of culture media for more various microorganisms are contained in the handbook "Manual of Methods for General Bacteriology" of the American Society for Bacteriology (Washington D.C., USA, 1981).

Sugars and carbohydrates, such as e.g. glucose, sucrose, lactose, fructose, maltose, molasses, starch and cellulose, oils and fats, such as e.g. soya oil, sunflower oil, groundnut oil and coconut fat, fatty acids, such as e.g.

palmitic acid, stearic acid and linoleic acid, alcohols, such as e.g. glycerol and ethanol, and organic acids, such as e.g. acetic acid, can be used as the source of carbon. These substances can be used individually or as a mixture.

- 5 Organic nitrogen-containing compounds, such as peptones, yeast extract, meat extract, malt extract, corn steep liquor, soya bean flour and urea, or inorganic compounds, such as ammonium sulfate, ammonium chloride, ammonium phosphate, ammonium carbonate and ammonium nitrate, can be
10 used as the source of nitrogen. The sources of nitrogen can be used individually or as a mixture.

- Potassium dihydrogen phosphate or dipotassium hydrogen phosphate or the corresponding sodium-containing salts can be used as the source of phosphorus. The culture medium
15 must furthermore comprise salts of metals, such as e.g. magnesium sulfate or iron sulfate, which are necessary for growth. Finally, essential growth substances, such as amino acids and vitamins, can be employed in addition to the abovementioned substances. Precursors of pantothenic acid,
20 such as aspartate, β -alanine, ketoisovalerate, ketopantoic acid or pantoic acid, and optionally salts thereof, can moreover be added to the culture medium to additionally increase the pantothenic acid production. The starting substances mentioned can be added to the culture in the
25 form of a single batch, or can be fed in during the culture in a suitable manner.

- Basic compounds, such as sodium hydroxide, potassium hydroxide, ammonia, or acid compounds, such as phosphoric acid or sulfuric acid, can be employed in a suitable manner
30 to control the pH of the culture. Antifoams, such as e.g. fatty acid polyglycol esters, can be employed to control the development of foam. Suitable substances having a selective action, e.g. antibiotics, can be added to the medium to maintain the stability of plasmids. To maintain
35 aerobic conditions, oxygen or oxygen-containing gas

mixtures, such as e.g. air, are introduced into the culture. The temperature of the culture is usually 20°C to 45°C, and preferably 25°C to 40°C. Culturing is continued until a maximum of pantothenic acid has formed. This target
5 is usually reached within 10 hours to 160 hours.

The concentration of pantothenic acid formed can be determined with known chemical (Velisek; Chromatographic Science 60, 515-560 (1992)) or microbiological methods, such as e.g. the *Lactobacillus plantarum* test (DIFCO
10 MANUAL, 10th Edition, p. 1100-1102; Michigan, USA).

The following microorganism was deposited on 19th October 1999 as a pure culture at the Deutsche Sammlung für Mikroorganismen und Zellkulturen (DSMZ = German Collection of Microorganisms and Cell Cultures, Braunschweig, Germany)
15 in accordance with the Budapest Treaty:

- *Escherichia coli* strain DH5 α /pCR2.1poxBint as DSM 13114.

The present invention is explained in more detail in the following with the aid of embodiment examples.

Example 1

20 Preparation of a genomic cosmid gene library from *Corynebacterium glutamicum* ATCC 13032

Chromosomal DNA from *C. glutamicum* ATCC 13032 was isolated as described by Tauch et al. (1995, Plasmid 33:168-179) and partly cleaved with the restriction enzyme *Sau3AI* (Amersham
25 Pharmacia, Freiburg, Germany, Product Description *Sau3AI*, Code no. 27-0913-02). The DNA fragments were dephosphorylated with shrimp alkaline phosphatase (Roche Molecular Biochemicals, Mannheim, Germany, Product Description SAP, Code no. 1758250). The DNA of the cosmid
30 vector SuperCos1 (Wahl et al. (1987) Proceedings of the National Academy of Sciences USA 84:2160-2164), obtained from Stratagene (La Jolla, USA, Product Description

SuperCos1 Cosmid Vector Kit, Code no. 251301) was cleaved with the restriction enzyme XbaI (Amersham Pharmacia, Freiburg, Germany, Product Description XbaI, Code no. 27-0948-02) and likewise dephosphorylated with shrimp alkaline phosphatase.

The cosmid DNA was then cleaved with the restriction enzyme BamHI (Amersham Pharmacia, Freiburg, Germany, Product Description BamHI, Code no. 27-0868-04). The cosmid DNA treated in this manner was mixed with the treated ATCC13032 DNA and the batch was treated with T4 DNA ligase (Amersham Pharmacia, Freiburg, Germany, Product Description T4-DNA-Ligase, Code no.27-0870-04). The ligation mixture was then packed in phages with the aid of Gigapack II XL Packing Extract (Stratagene, La Jolla, USA, Product Description Gigapack II XL Packing Extract, Code no. 200217). For infection of the E. coli strain NM554 (Raleigh et al. 1988, Nucleic Acid Res. 16:1563-1575) the cells were taken up in 10 mM MgSO₄ and mixed with an aliquot of the phage suspension. The infection and titering of the cosmid library were carried out as described by Sambrook et al. (1989, Molecular Cloning: A Laboratory Manual, Cold Spring Harbor), the cells being plated out on LB agar (Lennox, 1955, Virology, 1:190) with 100 µg/ml ampicillin. After incubation overnight at 37°C, recombinant individual clones were selected.

Example 2

Isolation and sequencing of the poxB gene

The cosmid DNA of an individual colony was isolated with the Qiaprep Spin Miniprep Kit (Product No. 27106, Qiagen, Hilden, Germany) in accordance with the manufacturer's instructions and partly cleaved with the restriction enzyme Sau3AI (Amersham Pharmacia, Freiburg, Germany, Product Description Sau3AI, Product No. 27-0913-02). The DNA fragments were dephosphorylated with shrimp alkaline

phosphatase (Roche Molecular Biochemicals, Mannheim, Germany, Product Description SAP, Product No. 1758250). After separation by gel electrophoresis, the cosmid fragments in the size range of 1500 to 2000 bp were
5 isolated with the QiaExII Gel Extraction Kit (Product No. 20021, Qiagen, Hilden, Germany).

The DNA of the sequencing vector pZero-1, obtained from Invitrogen (Groningen, The Netherlands, Product Description Zero Background Cloning Kit, Product No. K2500-01) was
10 cleaved with the restriction enzyme BamHI (Amersham Pharmacia, Freiburg, Germany, Product Description BamHI, Product No. 27-0868-04). The ligation of the cosmid fragments in the sequencing vector pZero-1 was carried out as described by Sambrook et al. (1989, Molecular Cloning: A
15 Laboratory Manual, Cold Spring Harbor), the DNA mixture being incubated overnight with T4 ligase (Pharmacia Biotech, Freiburg, Germany). This ligation mixture was then electroporated (Tauch et al., 1994, FEMS Microbiol Letters, 123:343-7) into the E. coli strain DH5 α MCR (Grant, 1990,
20 Proceedings of the National Academy of Sciences, U.S.A., 87:4645-4649) and plated out on LB agar (Lennox, 1955, Virology, 1:190) with 50 μ g/ml zeocin.

The plasmid preparation of the recombinant clones was carried out with the Biorobot 9600 (Product No. 900200,
25 Qiagen, Hilden, Germany). The sequencing was carried out by the dideoxy chain termination method of Sanger et al. (1977, Proceedings of the National Academies of Sciences, U.S.A., 74:5463-5467) with modifications according to Zimmermann et al. (1990, Nucleic Acids Research, 18:1067).
30 The "RR dRhodamin Terminator Cycle Sequencing Kit" from PE Applied Biosystems (Product No. 403044, Weiterstadt, Germany) was used. The separation by gel electrophoresis and analysis of the sequencing reaction were carried out in a "Rotiphoresis NF Acrylamide/Bisacrylamide" Gel (29:1)
35 (Product No. A124.1, Roth, Karlsruhe, Germany) with the

"ABI Prism 377" sequencer from PE Applied Biosystems (Weiterstadt, Germany).

The raw sequence data obtained were then processed using the Staden program package (1986, Nucleic Acids Research, 14:217-231) version 97-0. The individual sequences of the pZerol derivatives were assembled to a continuous contig. The computer-assisted coding region analysis were prepared with the XNIP program (Staden, 1986, Nucleic Acids Research, 14:217-231).

The resulting nucleotide sequence is shown in SEQ ID No. 1. Analysis of the nucleotide sequence showed an open reading frame of 1737 base pairs, which was called the poxB gene. The poxB gene codes for a polypeptide of 579 amino acids shown in SEQ ID No. 2.

Example 3

Preparation of the integration vector pCR2.1poxBint for mutagenesis of the poxB gene

From the strain ATCC 13032, chromosomal DNA was isolated by the method of Eikmanns et al. (Microbiology 140: 1817-1828 (1994)). On the basis of the sequence of the poxB gene known for *C. glutamicum* from example 2, the following oligonucleotides were chosen for the polymerase chain reaction:

poxBint1:

5' TGC GAG ATG GTG AAT GGT GG 3'

poxBint2:

5' GCA TGA GGC AAC GCA TTA GC 3'

The primers shown were synthesized by MWG Biotech (Ebersberg, Germany) and the PCR reaction was carried out by the standard PCR method of Innis et al. (PCR protocols. A Guide to Methods and Applications, 1990, Academic Press) with Pwo-Polymerase from Boehringer. With the aid of the

polymerase chain reaction, a DNA fragment approx. 0.9 kb in size was isolated, this carrying an internal fragment of the poxB gene and being shown in SEQ ID No. 3.

The amplified DNA fragment was ligated with the TOPO TA Cloning Kit from Invitrogen Corporation (Carlsbad, CA, USA; Catalogue Number K4500-01) in the vector pCR2.1-TOPO (Mead et al. (1991) Bio/Technology 9:657-663). The E. coli strain Top10F' (Grant et al. (1990) Proceedings of the National Academy of Sciences, USA, 87:4645-4649) was then electroporated. Selection for plasmid-carrying cells was made by plating out the transformation batch on LB agar (Sambrook et al., Molecular Cloning: A Laboratory Manual, 2nd Ed., Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y., 1989), which had been supplemented with 50 mg/l kanamycin. Plasmid DNA was isolated from a transformant with the aid of the QIAprep Spin Miniprep Kit from Qiagen and checked by restriction with the restriction enzyme EcoRI and subsequent agarose gel electrophoresis (0.8%). The plasmid was called pCR2.1poxBint (figure 1).

20 Example 4

Preparation of an exchange vector for deletion mutagenesis of the poxB gene

4.1 Determination of the nucleotide sequence of the flanks of the poxB gene

25 In further sequencing steps, the nucleotide sequence of the poxB gene region shown in SEQ ID No. 1 was extended upstream and downstream by in each case approx. 500 to 600 bp. The method described in example 2 was used for this. The extended nucleotide sequence of the poxB gene region shown in SEQ ID No. 4 was obtained in this manner. 30 The new upstream of the poxB gene region shown in SEQ ID No. 1 is shown in SEQ ID No. 6. The new downstream of the

poxB gene region shown in SEQ ID No. 1 is shown in SEQ ID No. 7.

4.2 Construction of a Δ poxB allele

The method of geneSOEing-PCR described by Horton (Molecular Microbiology 3:93-99 (1995)) was used for construction of the Δ poxB allele. The primer pairs shown in table 1 (see also SEQ ID No. 8 to 11) were constructed for this. By means of a PCR, the 5' region before the poxB gene was amplified with primer pair 1 and the 3' region after the poxB gene was amplified with primer pair 2. A further PCR was then carried out with the two amplification products and the primers pox-del1 and pox-del4, as a result of which the two amplification products were joined by means of geneSOEing. The deletion fragment or Δ poxB allele obtained in this way contains the flanking sequences of the poxB gene. The nucleotide sequence of the Δ poxB allele is shown in SEQ ID No. 12.

Table 1

Primer	5'-Sequence-3'	Primer pair
pox-del1	ATGAGGAACATCCGGCGGTG	1
pox-del2	GAGAACAGCAGGAGTATCAATCATCACTGAACT CCTCAACGTTATGGC	
pox-del3	TGATGATTGATACACCTGCTGTTCTC	2
pox-del4	TCATTGCCACCTGCTTCTCA	

4.3 Construction of an exchange vector

The DNA fragment obtained in this way was ligated with the Zero Blunt TOPO PCR Cloning Kit from Invitrogen Corporation (Carlsbad, CA, USA; Catalogue Number K2800-20) in the

vector pCR-Blunt II-TOPO vector (Shuman et al., (1994) Journal of Biological Chemistry 269:32678-32684; Bernard et al., (1983) Journal of Molecular Biology 234:534-541). The E. coli strain Top10 (Grant et al. (1990) Proceedings of the National Academy of Sciences, USA 87:4645-4649) was then transformed with the ligation batch. Selection for plasmid-carrying cells was made by plating out the transformation batch on LB agar (Sambrook et al., Molecular Cloning: A Laboratory Manual, 2nd Ed., Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y.), which had been supplemented with 50 mg/l kanamycin. Plasmid DNA was isolated from a transformant with the aid of the QIAprep Spin Miniprep Kit from Qiagen and checked by restriction with the restriction enzyme EcoRI and subsequent agarose gel electrophoresis (0.8%). The plasmid was called pCRB1-poxBdel (figure 2).

The insert carrying the Δ poxB allele was excised from this plasmid by means of EcoRI, isolated from the gel and ligated in the non-replicative integration vector pK18mobsacB, which was also cleaved with EcoRI (Schafer et al., Gene 145, 69-73 (1994)). The clonings were carried out in E. coli DH5 α mcr (Grant et al., (1990) Proceedings of the National Academy of Sciences, USA, 87: 4645-4649) as the host. The resulting plasmid was called pK18mobsacB-poxBdel.

The following figures are attached:

- Figure 1: Map of the plasmid pCR2.1poxBint
- Figure 2: Map of the plasmid pCRB1-poxBdel

The base pair numbers stated are approx. values obtained in the context of reproducibility.

The abbreviations and designations used have the following meaning:

Figure 1:

ApR	Ampicillin resistance gene
ColE1 ori	Replication origin ColE1
f1 ori	Replication origin of phage f1
KmR	Kanamycin resistance gene
lacZ	Residues of the lacZ α gene fragment
poxBint	Internal fragment of the poxB gene

Figure 2:

'lacZa	3' end of the lacZ α gene fragment
3'-Region	3' flank of the poxB gene
5'-Region	5' flank of the poxB gene
ccdB	ccdB gene
Km	Kanamycin resistance gene
lacZa'	5' end of the lacZ α gene fragment
plac	Promoter of the lac operon
pMB1	Replication origin of the plasmid pMB1
Zeocin	Zeocin resistance gene

The following abbreviations have moreover been used:

BamHI:	Cleavage site of the restriction enzyme BamHI
ClaI	Cleavage site of the restriction enzyme ClaI

EcoRI: Cleavage site of the restriction enzyme
EcoRI

HindIII: Cleavage site of the restriction enzyme
HindIII

SalI: Cleavage site of the restriction enzyme SalI

SEQUENCE PROTOCOL

<110> Degussa-Hüls AG

5 <120> Process for the fermentative preparation of
D-pantothenic acid using coryneform
bacteria.

10 <130> 000439 BT

<140>
<141>

<160> 12

15 <170> PatentIn Ver. 2.1

<210> 1
<211> 2160
20 <212> DNA
<213> Corynebacterium glutamicum

<220>
<221> CDS
25 <222> (327)..(2063)

<220>
<221> -35_signal
<222> (227)..(232)

30 <220>
<221> -10_signal
<222> (256)..(261)

35 <400> 1
ttagaggcga ttctgtgagg tcactttttg tggggtcggg gtctaaattt ggccagtttt 60
cgaggcgacc agacaggcgt gcccacgatg tttaaataagg cgatcgggtg gcatctgtgt 120
40 ttggtttcga cgggctgaaa ccaaaccaga ctgcccagca acgacggaaa tcccaaaagt 180
gggcatccct gtttggtacc gagtaccac ccgggcctga aactccctgg caggcggggcg 240
aagcgtggca acaactggaa tttaagagca caattgaagt cgcaccaagt taggcaacac 300
45 aatagccata acgttgagga gttcag atg gca cac agc tac gca gaa caa tta 353
Met Ala His Ser Tyr Ala Glu Gln Leu
1 5

50 att gac act ttg gaa gct caa ggt gtg aag cga att tat ggt ttg gtg 401
Ile Asp Thr Leu Glu Ala Gln Gly Val Lys Arg Ile Tyr Gly Leu Val
10 15 20 25

55 ggt gac agc ctt aat ccg atc gtg gat gct gtc cgc caa tca gat att 449
Gly Asp Ser Leu Asn Pro Ile Val Asp Ala Val Arg Gln Ser Asp Ile
30 35 40

		gag	tgg	gtg	cac	gtt	cga	aat	gag	gaa	gcg	gcg	gcg	ttt	gca	gcc	ggt	497
		Glu	Trp	Val	His	Val	Arg	Asn	Glu	Glu	Ala	Ala	Ala	Phe	Ala	Ala	Gly	
					45					50					55			
5		gcg	gaa	tcg	ttg	atc	act	ggg	gag	ctg	gca	gta	tgt	gct	gct	tct	tgt	545
		Ala	Glu	Ser	Leu	Ile	Thr	Gly	Glu	Leu	Ala	Val	Cys	Ala	Ala	Ser	Cys	
				60					65					70				
10		ggt	cct	gga	aac	aca	cac	ctg	att	cag	ggt	ctt	tat	gat	tcg	cat	cga	593
		Gly	Pro	Gly	Asn	Thr	His	Leu	Ile	Gln	Gly	Leu	Tyr	Asp	Ser	His	Arg	
			75					80					85					
15		aat	ggt	gcg	aag	gtg	ttg	gcc	atc	gct	agc	cat	att	ccg	agt	gcc	cag	641
		Asn	Gly	Ala	Lys	Val	Leu	Ala	Ile	Ala	Ser	His	Ile	Pro	Ser	Ala	Gln	
		90					95					100					105	
20		att	ggt	tcg	acg	ttc	ttc	cag	gaa	acg	cat	ccg	gag	att	ttg	ttt	aag	689
		Ile	Gly	Ser	Thr	Phe	Phe	Gln	Glu	Thr	His	Pro	Glu	Ile	Leu	Phe	Lys	
						110					115					120		
		gaa	tgc	tct	ggt	tac	tgc	gag	atg	gtg	aat	ggt	ggt	gag	cag	ggt	gaa	737
		Glu	Cys	Ser	Gly	Tyr	Cys	Glu	Met	Val	Asn	Gly	Gly	Glu	Gln	Gly	Glu	
					125					130					135			
25		cgc	att	ttg	cat	cac	gcg	att	cag	tcc	acc	atg	gcg	ggt	aaa	ggt	gtg	785
		Arg	Ile	Leu	His	His	Ala	Ile	Gln	Ser	Thr	Met	Ala	Gly	Lys	Gly	Val	
				140					145					150				
30		tcg	gtg	gta	gtg	att	cct	ggt	gat	atc	gct	aag	gaa	gac	gca	ggt	gac	833
		Ser	Val	Val	Val	Ile	Pro	Gly	Asp	Ile	Ala	Lys	Glu	Asp	Ala	Gly	Asp	
			155					160					165					
35		ggt	act	tat	tcc	aat	tcc	act	att	tct	tct	ggc	act	cct	gtg	gtg	ttc	881
		Gly	Thr	Tyr	Ser	Asn	Ser	Thr	Ile	Ser	Ser	Gly	Thr	Pro	Val	Val	Phe	
		170					175					180					185	
40		ccg	gat	cct	act	gag	gct	gca	gcg	ctg	gtg	gag	gcg	att	aac	aac	gct	929
		Pro	Asp	Pro	Thr	Glu	Ala	Ala	Ala	Leu	Val	Glu	Ala	Ile	Asn	Asn	Ala	
						190					195					200		
		aag	tct	gtc	act	ttg	ttc	tgc	ggt	gcg	ggc	gtg	aag	aat	gct	cgc	gcg	977
		Lys	Ser	Val	Thr	Leu	Phe	Cys	Gly	Ala	Gly	Val	Lys	Asn	Ala	Arg	Ala	
					205					210					215			
45		cag	gtg	ttg	gag	ttg	gcg	gag	aag	att	aaa	tca	ccg	atc	ggg	cat	gcg	1025
		Gln	Val	Leu	Glu	Leu	Ala	Glu	Lys	Ile	Lys	Ser	Pro	Ile	Gly	His	Ala	
				220					225					230				
50		ctg	ggt	ggt	aag	cag	tac	atc	cag	cat	gag	aat	ccg	ttt	gag	gtc	ggc	1073
		Leu	Gly	Gly	Lys	Gln	Tyr	Ile	Gln	His	Glu	Asn	Pro	Phe	Glu	Val	Gly	
			235					240					245					
55		atg	tct	ggc	ctg	ctt	ggt	tac	ggc	gcc	tgc	gtg	gat	gcg	tcc	aat	gag	1121
		Met	Ser	Gly	Leu	Leu	Gly	Tyr	Gly	Ala	Cys	Val	Asp	Ala	Ser	Asn	Glu	
		250					255					260					265	
		gcg	gat	ctg	ctg	att	cta	ttg	ggt	acg	gat	ttc	cct	tat	tct	gat	ttc	1169
		Ala	Asp	Leu	Leu	Ile	Leu	Leu	Gly	Thr	Asp	Phe	Pro	Tyr	Ser	Asp	Phe	
						270					275					280		

	ctt cct aaa gac aac gtt gcc cag gtg gat atc aac ggt gcg cac att	1217
	Leu Pro Lys Asp Asn Val Ala Gln Val Asp Ile Asn Gly Ala His Ile	
	285 290 295	
5	ggt cga cgt acc acg gtg aag tat ccg gtg acc ggt gat gtt gct gca	1265
	Gly Arg Arg Thr Thr Val Lys Tyr Pro Val Thr Gly Asp Val Ala Ala	
	300 305 310	
10	aca atc gaa aat att ttg cct cat gtg aag gaa aaa aca gat cgt tcc	1313
	Thr Ile Glu Asn Ile Leu Pro His Val Lys Glu Lys Thr Asp Arg Ser	
	315 320 325	
15	ttc ctt gat cgg atg ctc aag gca cac gag cgt aag ttg agc tcg gtg	1361
	Phe Leu Asp Arg Met Leu Lys Ala His Glu Arg Lys Leu Ser Ser Val	
	330 335 340 345	
20	gta gag acg tac aca cat aac gtc gag aag cat gtg cct att cac cct	1409
	Val Glu Thr Tyr Thr His Asn Val Glu Lys His Val Pro Ile His Pro	
	350 355 360	
25	gaa tac gtt gcc tct att ttg aac gag ctg gcg gat aag gat gcg gtg	1457
	Glu Tyr Val Ala Ser Ile Leu Asn Glu Leu Ala Asp Lys Asp Ala Val	
	365 370 375	
30	ttt act gtg gat acc ggc atg tgc aat gtg tgg cat gcg agg tac atc	1505
	Phe Thr Val Asp Thr Gly Met Cys Asn Val Trp His Ala Arg Tyr Ile	
	380 385 390	
35	gag aat ccg gag gga acg cgc gac ttt gtg ggt tca ttc cgc cac ggc	1553
	Glu Asn Pro Glu Gly Thr Arg Asp Phe Val Gly Ser Phe Arg His Gly	
	395 400 405	
40	acg atg gct aat gcg ttg cct cat gcg att ggt gcg caa agt gtt gat	1601
	Thr Met Ala Asn Ala Leu Pro His Ala Ile Gly Ala Gln Ser Val Asp	
	410 415 420 425	
45	cga aac cgc cag gtg atc gcg atg tgt ggc gat ggt ggt ttg ggc atg	1649
	Arg Asn Arg Gln Val Ile Ala Met Cys Gly Asp Gly Gly Leu Gly Met	
	430 435 440	
50	ctg ctg ggt gag ctt ctg acc gtt aag ctg cac caa ctt ccg ctg aag	1697
	Leu Leu Gly Glu Leu Leu Thr Val Lys Leu His Gln Leu Pro Leu Lys	
	445 450 455	
55	gct gtg gtg ttt aac aac agt tct ttg ggc atg gtg aag ttg gag atg	1745
	Ala Val Val Phe Asn Asn Ser Ser Leu Gly Met Val Lys Leu Glu Met	
	460 465 470	
60	ctc gtg gag gga cag cca gaa ttt ggt act gac cat gag gaa gtg aat	1793
	Leu Val Glu Gly Gln Pro Glu Phe Gly Thr Asp His Glu Glu Val Asn	
	475 480 485	
65	ttc gca gag att gcg gcg gct gcg ggt atc aaa tcg gta cgc atc acc	1841
	Phe Ala Glu Ile Ala Ala Ala Ala Gly Ile Lys Ser Val Arg Ile Thr	
	490 495 500 505	
70	gat ccg aag aaa gtt cgc gag cag cta gct gag gca ttg gca tat cct	1889
	Asp Pro Lys Lys Val Arg Glu Gln Leu Ala Glu Ala Leu Ala Tyr Pro	
	510 515 520	

gga cct gta ctg atc gat atc gtc acg gat cct aat gcg ctg tcg atc 1937
 Gly Pro Val Leu Ile Asp Ile Val Thr Asp Pro Asn Ala Leu Ser Ile
 525 530 535

5

cca cca acc atc acg tgg gaa cag gtc atg gga ttc agc aag gcg gcc 1985
 Pro Pro Thr Ile Thr Trp Glu Gln Val Met Gly Phe Ser Lys Ala Ala
 540 545 550

10

acc cga acc gtc ttt ggt gga gga gta gga gcg atg atc gat ctg gcc 2033
 Thr Arg Thr Val Phe Gly Gly Gly Val Gly Ala Met Ile Asp Leu Ala
 555 560 565

15

cgt tcg aac ata agg aat att cct act cca tgatgattga tacacctgct 2083
 Arg Ser Asn Ile Arg Asn Ile Pro Thr Pro
 570 575

gttctcattg accgcgagcg cttaactgcc aacatttcca ggatggcagc tcacgccggt 2143

20

gcccatgaga ttgccct 2160

<210> 2
 <211> 579
 <212> PRT
 <213> Corynebacterium glutamicum

25

<400> 2
 Met Ala His Ser Tyr Ala Glu Gln Leu Ile Asp Thr Leu Glu Ala Gln
 1 5 10 15

30

Gly Val Lys Arg Ile Tyr Gly Leu Val Gly Asp Ser Leu Asn Pro Ile
 20 25 30

35

Val Asp Ala Val Arg Gln Ser Asp Ile Glu Trp Val His Val Arg Asn
 35 40 45

Glu Glu Ala Ala Ala Phe Ala Ala Gly Ala Glu Ser Leu Ile Thr Gly
 50 55 60

40

Glu Leu Ala Val Cys Ala Ala Ser Cys Gly Pro Gly Asn Thr His Leu
 65 70 75 80

Ile Gln Gly Leu Tyr Asp Ser His Arg Asn Gly Ala Lys Val Leu Ala
 85 90 95

45

Ile Ala Ser His Ile Pro Ser Ala Gln Ile Gly Ser Thr Phe Phe Gln
 100 105 110

50

Glu Thr His Pro Glu Ile Leu Phe Lys Glu Cys Ser Gly Tyr Cys Glu
 115 120 125

Met Val Asn Gly Gly Glu Gln Gly Glu Arg Ile Leu His His Ala Ile
 130 135 140

55

Gln Ser Thr Met Ala Gly Lys Gly Val Ser Val Val Val Ile Pro Gly
 145 150 155 160

	Asp	Ile	Ala	Lys	Glu	Asp	Ala	Gly	Asp	Gly	Thr	Tyr	Ser	Asn	Ser	Thr	
					165					170					175		
5	Ile	Ser	Ser	Gly	Thr	Pro	Val	Val	Phe	Pro	Asp	Pro	Thr	Glu	Ala	Ala	
				180					185					190			
	Ala	Leu	Val	Glu	Ala	Ile	Asn	Asn	Ala	Lys	Ser	Val	Thr	Leu	Phe	Cys	
			195					200					205				
10	Gly	Ala	Gly	Val	Lys	Asn	Ala	Arg	Ala	Gln	Val	Leu	Glu	Leu	Ala	Glu	
		210					215					220					
	Lys	Ile	Lys	Ser	Pro	Ile	Gly	His	Ala	Leu	Gly	Gly	Lys	Gln	Tyr	Ile	
	225					230					235					240	
15	Gln	His	Glu	Asn	Pro	Phe	Glu	Val	Gly	Met	Ser	Gly	Leu	Leu	Gly	Tyr	
				245						250					255		
	Gly	Ala	Cys	Val	Asp	Ala	Ser	Asn	Glu	Ala	Asp	Leu	Leu	Ile	Leu	Leu	
20				260					265					270			
	Gly	Thr	Asp	Phe	Pro	Tyr	Ser	Asp	Phe	Leu	Pro	Lys	Asp	Asn	Val	Ala	
			275					280					285				
25	Gln	Val	Asp	Ile	Asn	Gly	Ala	His	Ile	Gly	Arg	Arg	Thr	Thr	Val	Lys	
		290					295					300					
	Tyr	Pro	Val	Thr	Gly	Asp	Val	Ala	Ala	Thr	Ile	Glu	Asn	Ile	Leu	Pro	
	305					310					315					320	
30	His	Val	Lys	Glu	Lys	Thr	Asp	Arg	Ser	Phe	Leu	Asp	Arg	Met	Leu	Lys	
				325						330					335		
	Ala	His	Glu	Arg	Lys	Leu	Ser	Ser	Val	Val	Glu	Thr	Tyr	Thr	His	Asn	
35				340					345					350			
	Val	Glu	Lys	His	Val	Pro	Ile	His	Pro	Glu	Tyr	Val	Ala	Ser	Ile	Leu	
			355					360					365				
40	Asn	Glu	Leu	Ala	Asp	Lys	Asp	Ala	Val	Phe	Thr	Val	Asp	Thr	Gly	Met	
		370					375					380					
	Cys	Asn	Val	Trp	His	Ala	Arg	Tyr	Ile	Glu	Asn	Pro	Glu	Gly	Thr	Arg	
	385					390					395					400	
45	Asp	Phe	Val	Gly	Ser	Phe	Arg	His	Gly	Thr	Met	Ala	Asn	Ala	Leu	Pro	
				405						410					415		
	His	Ala	Ile	Gly	Ala	Gln	Ser	Val	Asp	Arg	Asn	Arg	Gln	Val	Ile	Ala	
50				420					425					430			
	Met	Cys	Gly	Asp	Gly	Gly	Leu	Gly	Met	Leu	Leu	Gly	Glu	Leu	Leu	Thr	
			435					440					445				
55	Val	Lys	Leu	His	Gln	Leu	Pro	Leu	Lys	Ala	Val	Val	Phe	Asn	Asn	Ser	
		450					455					460					
	Ser	Leu	Gly	Met	Val	Lys	Leu	Glu	Met	Leu	Val	Glu	Gly	Gln	Pro	Glu	
	465					470					475					480	

Phe Gly Thr Asp His Glu Glu Val Asn Phe Ala Glu Ile Ala Ala Ala
 485 490 495
 5 Ala Gly Ile Lys Ser Val Arg Ile Thr Asp Pro Lys Lys Val Arg Glu
 500 505 510
 Gln Leu Ala Glu Ala Leu Ala Tyr Pro Gly Pro Val Leu Ile Asp Ile
 515 520 525
 10 Val Thr Asp Pro Asn Ala Leu Ser Ile Pro Pro Thr Ile Thr Trp Glu
 530 535 540
 Gln Val Met Gly Phe Ser Lys Ala Ala Thr Arg Thr Val Phe Gly Gly
 15 545 550 555 560
 Gly Val Gly Ala Met Ile Asp Leu Ala Arg Ser Asn Ile Arg Asn Ile
 565 570 575
 20 Pro Thr Pro
 25 <210> 3
 <211> 875
 <212> DNA
 <213> *Corynebacterium glutamicum*
 30 <400> 3
 tgcgagatgg tgaatggtgg tgagcagggt gaacgcattt tgcatacagc gattcagttcc 60
 accatggcgg gtaaagggtg gtcgggtgga gtgattcctg gtgatatcgc taaggaagac 120
 gcaggtgacg gtacttattc caattccact atttcttctg gcactcctgt ggtgttcccg 180
 gatcctactg aggctgcagc gctggtggag gcgattaaca acgctaagtc tgtcactttg 240
 35 ttctgcggtg cgggcgtgaa gaatgctcgc gcgcagggtg tggagttggc ggagaagatt 300
 aatcaccga tcgggcatgc gctgggtggt aagcagtaca tccagcatga gaatccggtt 360
 gaggtcggca tgtctggcct gcttggttac ggcgcctgcg tggatgcgtc caatgaggcg 420
 gatctgctga ttctattggg tacggatttc ccttattctg atttccttcc taaagacaac 480
 gttgcccgag tggatatcaa cgggtgcgcac attggtcgac gtaccacggg gaagtatccg 540
 40 gtgaccggtg atgttgctgc aacaatcgaa aatattttgc ctcatgtgaa ggaaaaaaca 600
 gatcgttcct tccttgatcg gatgctcaag gcacacgagc gtaagttgag ctcggtggta 660
 gagacgtaca cacataacgt cgagaagcat gtgcctattc accctgaata cgttgccctc 720
 attttgaacg agctggcgga taaggatgcg gtgtttactg tggataaccg catgtgcaat 780
 gtgtggcatg cgaggtacat cgagaatccg gagggaaacg gcgactttgt gggttcattc 840
 45 cgccacggca cgatggctaa tgcgttgccct catgc 875
 <210> 4
 <211> 3248
 50 <212> DNA
 <213> *Corynebacterium glutamicum*
 <220>
 <221> CDS
 55 <222> (802)..(2538)
 <400> 4
 gctctcgag caacaagagc ccacgcagtt ggagcaaacg cagcaccaag tgaagcgatt 60

	ccgaaaatgc	tcaagcccat	gaggaacatc	cggcgggtggc	cgattttgtc	acccaaagtg	120										
	ccggtaccca	aaagaaggcc	cgccatgagc	aggggatatg	cgttgatgat	ccacaacgct	180										
5	tgggtttcgg	tggctgcgag	ctgttcacgc	agcagaggga	gtgcgggtgta	gagaatcgag	240										
	ttgtctacac	cgatcagaaa	gagaccaccg	ctgataacgg	cgaggaaagc	ccaacgtttg	300										
10	gttttcgtag	gcgcttgcg	ctgtaagggt	tctgaagtca	tggatcgtaa	ctgtaacgaa	360										
	tggtcgggtac	agttacaact	cttttggttg	tgttttagac	cacggcgctg	tgtggcgatt	420										
	taagacgtcg	gaaatcgtag	gggactgtca	gcgtgggtcg	ggttctttga	ggcgcttaga	480										
15	ggcgattctg	tgaggtcact	ttttgtgggg	tcgggggtcta	aatttggcc	gttttcgagg	540										
	cgaccagaca	ggcgtgccca	cgatgtttta	ataggcgatc	ggtagggcatc	tgtgttttgg	600										
20	ttcgacgggc	tgaacacaaa	ccagactgcc	cagcaacgac	ggaaatccca	aaagtgggca	660										
	tccctgtttg	gtaccgagta	cccaccggg	cctgaaactc	cctggcaggc	gggcgaagcg	720										
	tggcaacaac	tggaatttaa	gagcacaatt	gaagtcgcac	caagttaggc	aacacaatag	780										
25	ccataacggt	gaggagtcca	g	atg	gca	cac	agc	tac	gca	gaa	caa	tta	att		831		
				Met	Ala	His	Ser	Tyr	Ala	Glu	Gln	Leu	Ile				
				1				5					10				
30	gac	act	ttg	gaa	gct	caa	ggg	gtg	aag	cga	att	tat	ggg	ttg	gtg	ggg	879
	Asp	Thr	Leu	Glu	Ala	Gln	Gly	Val	Lys	Arg	Ile	Tyr	Gly	Leu	Val	Gly	
					15					20					25		
	gac	agc	ctt	aat	ccg	atc	gtg	gat	gct	gtc	cgc	caa	tca	gat	att	gag	927
35	Asp	Ser	Leu	Asn	Pro	Ile	Val	Asp	Ala	Val	Arg	Gln	Ser	Asp	Ile	Glu	
				30					35					40			
	tgg	gtg	cac	gtt	cga	aat	gag	gaa	gcg	gcg	gcg	ttt	gca	gcc	ggg	gcg	975
40	Trp	Val	His	Val	Arg	Asn	Glu	Glu	Ala	Ala	Ala	Phe	Ala	Ala	Gly	Ala	
			45					50					55				
	gaa	tcg	ttg	atc	act	ggg	gag	ctg	gca	gta	tgt	gct	gct	tct	tgt	ggg	1023
	Glu	Ser	Leu	Ile	Thr	Gly	Glu	Leu	Ala	Val	Cys	Ala	Ala	Ser	Cys	Gly	
		60					65					70					
45	cct	gga	aac	aca	cac	ctg	att	cag	ggg	ctt	tat	gat	tcg	cat	cga	aat	1071
	Pro	Gly	Asn	Thr	His	Leu	Ile	Gln	Gly	Leu	Tyr	Asp	Ser	His	Arg	Asn	
		75				80					85					90	
50	ggg	gcg	aag	gtg	ttg	gcc	atc	gct	agc	cat	att	ccg	agt	gcc	cag	att	1119
	Gly	Ala	Lys	Val	Leu	Ala	Ile	Ala	Ser	His	Ile	Pro	Ser	Ala	Gln	Ile	
					95				100						105		
	ggg	tcg	acg	ttc	ttc	cag	gaa	acg	cat	ccg	gag	att	ttg	ttt	aag	gaa	1167
55	Gly	Ser	Thr	Phe	Phe	Gln	Glu	Thr	His	Pro	Glu	Ile	Leu	Phe	Lys	Glu	
				110					115					120			
	tgc	tct	ggg	tac	tgc	gag	atg	gtg	aat	ggg	ggg	gag	cag	ggg	gaa	cgc	1215
	Cys	Ser	Gly	Tyr	Cys	Glu	Met	Val	Asn	Gly	Gly	Glu	Gln	Gly	Glu	Arg	
			125					130					135				

	att	ttg	cat	cac	gcg	att	cag	tcc	acc	atg	gcg	ggt	aaa	ggt	gtg	tcg	1263
	Ile	Leu	His	His	Ala	Ile	Gln	Ser	Thr	Met	Ala	Gly	Lys	Gly	Val	Ser	
		140					145					150					
5	gtg	gta	gtg	att	cct	ggt	gat	atc	gct	aag	gaa	gac	gca	ggt	gac	ggt	1311
	Val	Val	Val	Ile	Pro	Gly	Asp	Ile	Ala	Lys	Glu	Asp	Ala	Gly	Asp	Gly	
		155				160					165					170	
10	act	tat	tcc	aat	tcc	act	att	tct	tct	ggc	act	cct	gtg	gtg	ttc	ccg	1359
	Thr	Tyr	Ser	Asn	Ser	Thr	Ile	Ser	Ser	Gly	Thr	Pro	Val	Val	Phe	Pro	
					175					180					185		
15	gat	cct	act	gag	gct	gca	gcg	ctg	gtg	gag	gcg	att	aac	aac	gct	aag	1407
	Asp	Pro	Thr	Glu	Ala	Ala	Ala	Leu	Val	Glu	Ala	Ile	Asn	Asn	Ala	Lys	
				190					195					200			
20	tct	gtc	act	ttg	ttc	tgc	ggt	gcg	ggc	gtg	aag	aat	gct	cgc	gcg	cag	1455
	Ser	Val	Thr	Leu	Phe	Cys	Gly	Ala	Gly	Val	Lys	Asn	Ala	Arg	Ala	Gln	
			205					210					215				
25	gtg	ttg	gag	ttg	gcg	gag	aag	att	aaa	tca	ccg	atc	ggg	cat	gcg	ctg	1503
	Val	Leu	Glu	Leu	Ala	Glu	Lys	Ile	Lys	Ser	Pro	Ile	Gly	His	Ala	Leu	
		220					225				230						
30	ggt	ggt	aag	cag	tac	atc	cag	cat	gag	aat	ccg	ttt	gag	gtc	ggc	atg	1551
	Gly	Gly	Lys	Gln	Tyr	Ile	Gln	His	Glu	Asn	Pro	Phe	Glu	Val	Gly	Met	
		235				240					245					250	
35	tct	ggc	ctg	ctt	ggt	tac	ggc	gcc	tgc	gtg	gat	gcg	tcc	aat	gag	gcg	1599
	Ser	Gly	Leu	Leu	Gly	Tyr	Gly	Ala	Cys	Val	Asp	Ala	Ser	Asn	Glu	Ala	
					255				260						265		
40	gat	ctg	ctg	att	cta	ttg	ggt	acg	gat	ttc	cct	tat	tct	gat	ttc	ctt	1647
	Asp	Leu	Leu	Ile	Leu	Leu	Gly	Thr	Asp	Phe	Pro	Tyr	Ser	Asp	Phe	Leu	
				270					275					280			
45	cct	aaa	gac	aac	gtt	gcc	cag	gtg	gat	atc	aac	ggt	gcg	cac	att	ggt	1695
	Pro	Lys	Asp	Asn	Val	Ala	Gln	Val	Asp	Ile	Asn	Gly	Ala	His	Ile	Gly	
			285					290					295				
50	cga	cgt	acc	acg	gtg	aag	tat	ccg	gtg	acc	ggt	gat	gtt	gct	gca	aca	1743
	Arg	Arg	Thr	Thr	Val	Lys	Tyr	Pro	Val	Thr	Gly	Asp	Val	Ala	Ala	Thr	
		300					305					310					
55	atc	gaa	aat	att	ttg	cct	cat	gtg	aag	gaa	aaa	aca	gat	cgt	tcc	ttc	1791
	Ile	Glu	Asn	Ile	Leu	Pro	His	Val	Lys	Glu	Lys	Thr	Asp	Arg	Ser	Phe	
		315				320					325					330	
60	ctt	gat	cgg	atg	ctc	aag	gca	cac	gag	cgt	aag	ttg	agc	tcg	gtg	gta	1839
	Leu	Asp	Arg	Met	Leu	Lys	Ala	His	Glu	Arg	Lys	Leu	Ser	Ser	Val	Val	
					335					340					345		
65	gag	acg	tac	aca	cat	aac	gtc	gag	aag	cat	gtg	cct	att	cac	cct	gaa	1887
	Glu	Thr	Tyr	Thr	His	Asn	Val	Glu	Lys	His	Val	Pro	Ile	His	Pro	Glu	
				350					355					360			
70	tac	gtt	gcc	tct	att	ttg	aac	gag	ctg	gcg	gat	aag	gat	gcg	gtg	ttt	1935
	Tyr	Val	Ala	Ser	Ile	Leu	Asn	Glu	Leu	Ala	Asp	Lys	Asp	Ala	Val	Phe	
			365					370					375				

5	act	gtg	gat	acc	ggc	atg	tgc	aat	gtg	tgg	cat	gcg	agg	tac	atc	gag	1983		
	Thr	Val	Asp	Thr	Gly	Met	Cys	Asn	Val	Trp	His	Ala	Arg	Tyr	Ile	Glu			
	380						385					390							
	aat	ccg	gag	gga	acg	cgc	gac	ttt	gtg	ggt	tca	ttc	cgc	cac	ggc	acg	2031		
	Asn	Pro	Glu	Gly	Thr	Arg	Asp	Phe	Val	Gly	Ser	Phe	Arg	His	Gly	Thr			
	395					400					405					410			
10	atg	gct	aat	gcg	ttg	cct	cat	gcg	att	ggt	gcg	caa	agt	gtt	gat	cga	2079		
	Met	Ala	Asn	Ala	Leu	Pro	His	Ala	Ile	Gly	Ala	Gln	Ser	Val	Asp	Arg			
					415					420					425				
15	aac	cgc	cag	gtg	atc	gcg	atg	tgt	ggc	gat	ggt	ggt	ttg	ggc	atg	ctg	2127		
	Asn	Arg	Gln	Val	Ile	Ala	Met	Cys	Gly	Asp	Gly	Gly	Leu	Gly	Met	Leu			
					430					435					440				
20	ctg	ggt	gag	ctt	ctg	acc	gtt	aag	ctg	cac	caa	ctt	ccg	ctg	aag	gct	2175		
	Leu	Gly	Glu	Leu	Leu	Thr	Val	Lys	Leu	His	Gln	Leu	Pro	Leu	Lys	Ala			
					445					450					455				
25	gtg	gtg	ttt	aac	aac	agt	tct	ttg	ggc	atg	gtg	aag	ttg	gag	atg	ctc	2223		
	Val	Val	Phe	Asn	Asn	Ser	Ser	Leu	Gly	Met	Val	Lys	Leu	Glu	Met	Leu			
						460					465					470			
	gtg	gag	gga	cag	cca	gaa	ttt	ggt	act	gac	cat	gag	gaa	gtg	aat	ttc	2271		
	Val	Glu	Gly	Gln	Pro	Glu	Phe	Gly	Thr	Asp	His	Glu	Glu	Val	Asn	Phe			
						475					480					485			
30	gca	gag	att	gcg	gcg	gct	gcg	ggt	atc	aaa	tcg	gta	cgc	atc	acc	gat	2319		
	Ala	Glu	Ile	Ala	Ala	Ala	Ala	Gly	Ile	Lys	Ser	Val	Arg	Ile	Thr	Asp			
					495					500					505				
35	ccg	aag	aaa	gtt	cgc	gag	cag	cta	gct	gag	gca	ttg	gca	tat	cct	gga	2367		
	Pro	Lys	Lys	Val	Arg	Glu	Gln	Leu	Ala	Glu	Ala	Leu	Ala	Tyr	Pro	Gly			
					510					515					520				
40	cct	gta	ctg	atc	gat	atc	gtc	acg	gat	cct	aat	gcg	ctg	tcg	atc	cca	2415		
	Pro	Val	Leu	Ile	Asp	Ile	Val	Thr	Asp	Pro	Asn	Ala	Leu	Ser	Ile	Pro			
					525					530					535				
45	cca	acc	atc	acg	tgg	gaa	cag	gtc	atg	gga	ttc	agc	aag	gcg	gcc	acc	2463		
	Pro	Thr	Ile	Thr	Trp	Glu	Gln	Val	Met	Gly	Phe	Ser	Lys	Ala	Ala	Thr			
						540					545					550			
	cga	acc	gtc	ttt	ggt	gga	gga	gta	gga	gcg	atg	atc	gat	ctg	gcc	cgt	2511		
	Arg	Thr	Val	Phe	Gly	Gly	Gly	Val	Gly	Ala	Met	Ile	Asp	Leu	Ala	Arg			
						555					560					565			
50	tcg	aac	ata	agg	aat	att	cct	act	cca	tgatgattga	tacacctgct						2558		
	Ser	Asn	Ile	Arg	Asn	Ile	Pro	Thr	Pro										
					575														
55	gtttctcattg			accgcgagcg			cttaactgcc			aacattttcca			ggatggcagc			tcacgccggt			2618
	gcccatgaga			ttgccctgcg			tccgcatgtg			aaaacgcaca			aaatcattga			aattgcgcag			2678

tttgccggcg caggttttac ggacatcttt attgcatatc cgctgtatct aaccgatcat 2798
 5 gcaagtgaac gcctgaacgc gatccccgga gaaatttcca ttggcgtgga ttcggtagag 2858
 atggcacagg cgacggcgggg tttgcgggaa gatatcaagg ctctgattga agtggattcg 2918
 ggacatcgta gaagtggagt cacggcgact gcttcagaat tgagtcagat ccgcgaggcg 2978
 10 ctgggcagca ggtatgcagg agtggtttact tttcctgggc attcttatgg cccgggaaat 3038
 ggtgagcagg cagcagctga tgagcttcag gctctaaaca acagcgtcca gcgacttgct 3098
 15 ggcggcctga cttctggcgg ttcctcgccg tctgcgcagt ttacagacgc aatcgatgag 3158
 atgcgaccag gcgtgtatgt gtttaacgat tcccagcaga tcacctcggg agcatgcact 3218
 gagaagcagg tggcaatgac ggtgctgtct 3248
 20
 <210> 5
 <211> 579
 <212> PRT
 <213> *Corynebacterium glutamicum*
 25
 <400> 5
 Met Ala His Ser Tyr Ala Glu Gln Leu Ile Asp Thr Leu Glu Ala Gln
 1 5 10 15
 30 Gly Val Lys Arg Ile Tyr Gly Leu Val Gly Asp Ser Leu Asn Pro Ile
 20 25 30
 Val Asp Ala Val Arg Gln Ser Asp Ile Glu Trp Val His Val Arg Asn
 35 35 40 45
 Glu Glu Ala Ala Ala Phe Ala Ala Gly Ala Glu Ser Leu Ile Thr Gly
 50 55 60
 40 Glu Leu Ala Val Cys Ala Ala Ser Cys Gly Pro Gly Asn Thr His Leu
 65 70 75 80
 Ile Gln Gly Leu Tyr Asp Ser His Arg Asn Gly Ala Lys Val Leu Ala
 85 90 95
 45 Ile Ala Ser His Ile Pro Ser Ala Gln Ile Gly Ser Thr Phe Phe Gln
 100 105 110
 Glu Thr His Pro Glu Ile Leu Phe Lys Glu Cys Ser Gly Tyr Cys Glu
 115 120 125
 50 Met Val Asn Gly Gly Glu Gln Gly Glu Arg Ile Leu His His Ala Ile
 130 135 140
 55 Gln Ser Thr Met Ala Gly Lys Gly Val Ser Val Val Val Ile Pro Gly
 145 150 155 160
 Asp Ile Ala Lys Glu Asp Ala Gly Asp Gly Thr Tyr Ser Asn Ser Thr
 165 170 175

	Ile	Ser	Ser	Gly	Thr	Pro	Val	Val	Phe	Pro	Asp	Pro	Thr	Glu	Ala	Ala	
				180					185					190			
5	Ala	Leu	Val	Glu	Ala	Ile	Asn	Asn	Ala	Lys	Ser	Val	Thr	Leu	Phe	Cys	
			195					200					205				
	Gly	Ala	Gly	Val	Lys	Asn	Ala	Arg	Ala	Gln	Val	Leu	Glu	Leu	Ala	Glu	
		210					215					220					
10	Lys	Ile	Lys	Ser	Pro	Ile	Gly	His	Ala	Leu	Gly	Gly	Lys	Gln	Tyr	Ile	
	225					230					235					240	
	Gln	His	Glu	Asn	Pro	Phe	Glu	Val	Gly	Met	Ser	Gly	Leu	Leu	Gly	Tyr	
				245						250					255		
15	Gly	Ala	Cys	Val	Asp	Ala	Ser	Asn	Glu	Ala	Asp	Leu	Leu	Ile	Leu	Leu	
				260					265					270			
20	Gly	Thr	Asp	Phe	Pro	Tyr	Ser	Asp	Phe	Leu	Pro	Lys	Asp	Asn	Val	Ala	
			275					280					285				
	Gln	Val	Asp	Ile	Asn	Gly	Ala	His	Ile	Gly	Arg	Arg	Thr	Thr	Val	Lys	
		290					295					300					
25	Tyr	Pro	Val	Thr	Gly	Asp	Val	Ala	Ala	Thr	Ile	Glu	Asn	Ile	Leu	Pro	
	305					310					315					320	
	His	Val	Lys	Glu	Lys	Thr	Asp	Arg	Ser	Phe	Leu	Asp	Arg	Met	Leu	Lys	
					325					330				335			
30	Ala	His	Glu	Arg	Lys	Leu	Ser	Ser	Val	Val	Glu	Thr	Tyr	Thr	His	Asn	
				340					345					350			
	Val	Glu	Lys	His	Val	Pro	Ile	His	Pro	Glu	Tyr	Val	Ala	Ser	Ile	Leu	
35			355				360						365				
	Asn	Glu	Leu	Ala	Asp	Lys	Asp	Ala	Val	Phe	Thr	Val	Asp	Thr	Gly	Met	
		370					375					380					
40	Cys	Asn	Val	Trp	His	Ala	Arg	Tyr	Ile	Glu	Asn	Pro	Glu	Gly	Thr	Arg	
	385					390					395					400	
	Asp	Phe	Val	Gly	Ser	Phe	Arg	His	Gly	Thr	Met	Ala	Asn	Ala	Leu	Pro	
				405						410					415		
45	His	Ala	Ile	Gly	Ala	Gln	Ser	Val	Asp	Arg	Asn	Arg	Gln	Val	Ile	Ala	
				420					425					430			
	Met	Cys	Gly	Asp	Gly	Gly	Leu	Gly	Met	Leu	Leu	Gly	Glu	Leu	Leu	Thr	
50			435					440					445				
	Val	Lys	Leu	His	Gln	Leu	Pro	Leu	Lys	Ala	Val	Val	Phe	Asn	Asn	Ser	
		450					455					460					
55	Ser	Leu	Gly	Met	Val	Lys	Leu	Glu	Met	Leu	Val	Glu	Gly	Gln	Pro	Glu	
	465					470					475					480	
	Phe	Gly	Thr	Asp	His	Glu	Glu	Val	Asn	Phe	Ala	Glu	Ile	Ala	Ala	Ala	
					485					490						495	

Ala Gly Ile Lys Ser Val Arg Ile Thr Asp Pro Lys Lys Val Arg Glu
500 505 510

5 Gln Leu Ala Glu Ala Leu Ala Tyr Pro Gly Pro Val Leu Ile Asp Ile
515 520 525

Val Thr Asp Pro Asn Ala Leu Ser Ile Pro Pro Thr Ile Thr Trp Glu
530 535 540

10 Gln Val Met Gly Phe Ser Lys Ala Ala Thr Arg Thr Val Phe Gly Gly
545 550 555 560

15 Gly Val Gly Ala Met Ile Asp Leu Ala Arg Ser Asn Ile Arg Asn Ile
565 570 575

Pro Thr Pro

20

<210> 6
<211> 475
<212> DNA
25 <213> *Corynebacterium glutamicum*

<400> 6
gctctcgcag caacaagagc ccacgcagtt ggagcaaacg cagcaccaag tgaagcgatt 60
ccgaaaatgc tcaagcccat gaggaacatc cggcgggtggc cgattttgtc acccaaagtg 120
30 ccggtaccca aaagaaggcc cgccatgagc aggggatatg cgttgatgat ccacaacgct 180
tgggtttcgg tggctgcgag ctgttcacgc agcagaggga gtgcggtgta gagaatcgag 240
ttgtctacac cgatcagaaa gagaccaccg ctgataacgg cgaggaaaagc ccaacgttgg 300
gttttcgtag gcgcttgccg ctgtaagggt tctgaagtca tggatcgtaa ctgtaacgaa 360
tggtcggtag agttacaact cttttgttgg tgttttagac cacggcgctg tgtggcgatt 420
35 taagacgtcg gaaatcgtag gggactgtca gcgtgggtcg ggttctttga ggcgc 475

<210> 7
<211> 613
40 <212> DNA
<213> *Corynebacterium glutamicum*

<400> 7
gcgtccgcat gtgaaaacgc acaaaatcat tgaaattgcg cagatgcagg tcgacgccgg 60
45 tgcccagagg atcacctgcg caaccattgg cgaggcggaa atttttgccg gcgcagggtt 120
tacggacatc tttattgcat atccgctgta tctaaccgat catgcagtgc aacgcctgaa 180
cgcgatcccc ggagaaattt ccattggcgt ggattcggtg gagatggcac aggcgacggc 240
gggtttgcgg gaagatatca aggctctgat tgaagtggat tcgggacatc gtagaagtgg 300
agtcacggcg actgcttcag aattgagtca gatccgcgag gcgctgggca gcaggatatgc 360
50 aggagtgttt acttttctg ggcattctta tggcccggga aatggtgagc aggcagcagc 420
tgatgagctt caggctctaa acaacagcgt ccagcgactt gctggcggcc tgacttctgg 480
cggttctctg ccgtctgcgc agtttacaga cgcaatcgat gagatgcgac caggcgtgta 540
tgtgtttaac gattcccagc agatcacctc gggagcatgc actgagaagc aggtggcaat 600
55 gacgggtgctg tct 613

<210> 8
<211> 20
<212> DNA

<213> Artificial sequence
 <220>
 <223> Description of the artificial sequence: Primer
 5 pox-del1
 <400> 8
 atgaggaaca tccggcggtg 20
 10
 <210> 9
 <211> 48
 <212> DNA
 <213> Artificial sequence
 15
 <220>
 <223> Description of the artificial sequence: Primer
 pox-del2
 20 <400> 9
 gagaacagca ggagtatcaa tcatcactga actcctcaac gttatggc 48
 25
 <210> 10
 <211> 26
 <212> DNA
 <213> Artificial sequence
 <220>
 30 <223> Description of the artificial sequence: Primer
 pox-del3
 <400> 10
 35 tgatgattga tacacctgct gttctc 26
 <210> 11
 <211> 20
 <212> DNA
 40 <213> Artificial sequence
 <220>
 <223> Description of the artificial sequence: Primer
 pox-del4
 45 <400> 11
 tcattgccac ctgcttctca 20
 50
 <210> 12
 <211> 1422
 <212> DNA
 <213> Corynebacterium glutamicum
 55
 <220>
 <221> misc_feature
 <222> (1)..(1422)
 <223> Sequence of the delta poxB allele

<220>

<221> misc_feature

<222> (723)..(724)

Deletion of the coding region of the poxB gene

5

<400> 12

	atgaggaaca	tccggcgggtg	gccgattttg	tcacccaaag	tgccggtacc	caaaagaagg	60
	cccgccatga	gcaggggata	tgcgttgatg	atccacaacg	cttgggtttc	ggtggctgcg	120
	agctgttcac	gcagcagagg	gagtgcgggtg	tagagaatcg	agttgtctac	accgatcaga	180
10	aagagaccac	cgctgataac	ggcgaggaaa	gcccacggtt	gggttttcgt	aggcgcttgc	240
	gcctgtaagg	tttctgaagt	catggatcgt	aactgtaacg	aatggtcggt	acagttacaa	300
	ctcttttgtt	ggtgttttag	accacggcgc	tgtgtggcga	tttaagacgt	cggaaatcgt	360
	aggggactgt	cagcgtgggt	cgggttcttt	gaggcgctta	gaggcgattc	tgtgagggtc	420
	ctttttgtgg	ggtcggggtc	taaatttggc	cagttttcga	ggcgaccaga	caggcggtgcc	480
15	cacgatgttt	aaataggcga	tcggtgggca	tctgtgtttg	gtttcgacgg	gctgaaacca	540
	aaccagactg	cccagcaacg	acggaaatcc	caaaagtggg	catccctggt	tggtaccgag	600
	taccaccccg	ggcctgaaac	tccctggcag	gcgggcgaag	cgtggcaaca	actggaattt	660
	aagagcacia	ttgaagtgcg	accaagttag	gcaacacaat	agccataaag	ttgaggagtt	720
	cagtgatgat	tgatacacct	gctgttctca	ttgaccgcga	gcgcttaact	gccaacatth	780
20	ccaggatggc	agctcacgcc	ggtgcccattg	agattgccct	gcgtccgcgt	gtgaaaacgc	840
	acaaaatcat	tgaaattgcg	cagatgcagg	tcgacgcccg	tgcccagagg	atcacctgcg	900
	caaccattgg	cgaggcggaa	atthtttgccg	gcgcaggttt	tacggacatc	tttattgcat	960
	atccgctgta	tctaaccgat	catgcagtgc	aacgcctgaa	cgcgatcccc	ggagaaaatth	1020
	ccattggcgt	ggattcggta	gagatggcac	aggcgacggc	gggtttgctg	gaagatatca	1080
25	aggctctgat	tgaagtggat	tcgggacatc	gtagaagtgg	agtcacggcg	actgcttcag	1140
	aattgagtca	gatccgcgag	gcgctgggca	gcaggtatgc	aggagtgttt	actthttcctg	1200
	ggcattctta	tggcccggga	aatggtgagc	aggcagcagc	tgatgagctt	caggctctaa	1260
	acaacagcgt	ccagcgactt	gctggcggcc	tgacttctgg	cggttcctcg	ccgtctgcgc	1320
	agtttacaga	cgcaatcgat	gagatgcgac	caggcgtgta	tgtgtttaac	gattcccagc	1380
30	agatcacctc	gggagcatgc	actgagaagc	aggtggcaat	ga		1422

Patent claims

1. A process for the fermentative preparation of D-pantothenic acid, wherein the following steps are carried out:
 - 5 a) fermentation of D-pantothenic acid-producing coryneform bacteria in which at least the nucleotide sequence which codes for pyruvate oxidase (EC 1.2.2.2) (poxB) is attenuated, in particular eliminated;
 - 10 b) concentration of the D-pantothenic acid in the medium or in the cells of the bacteria; and
 - c) isolation of the D-pantothenic acid produced.
2. The process as claimed in claim 1, wherein the process
15 of insertion, in particular by means of the vector pCR2.1poxBint, shown in figure 1 and deposited in E.coli as DSM 13114, is used to achieve the attenuation.
3. The process as claimed in claim 1, wherein the process
20 of deletion, in particular by means of the vector pCRB1-poxBdel, shown in figure 2, is used to achieve the attenuation.
4. The process as claimed in claim 1, wherein coryneform
25 bacteria in which further genes of the biosynthesis pathway of D-pantothenic acid are additionally enhanced are employed.
5. The process as claimed in claim 1, wherein coryneform
30 bacteria in which the metabolic pathways which reduce the formation of D-pantothenic acid are at least partly eliminated are employed.
6. The process as claimed in claim 3, wherein coryneform
bacteria in which at the same time the panB gene which

codes for ketopantoate hydroxymethyl transferase is enhanced are employed.

- 5 7. The process as claimed in claim 3, wherein coryneform bacteria in which at the same time the panC gene which codes for pantothenate is enhanced are employed.
8. The process as claimed in claim 3, wherein coryneform bacteria in which at the same time the ilvC gene which codes for acetohydroxy-acid isomeroreductase is enhanced are employed.
- 10 9. The process as claimed in claim 3, wherein coryneform bacteria in which at the same time the ilvD gene which codes for dihydroxy-acid dehydratase is enhanced are employed.
- 15 10. The process as claimed in claim 1 or 4, wherein coryneform bacteria in which at the same time the pck gene which codes for phosphoenol pyruvate carboxykinase is attenuated are employed.
- 20 11. The process as claimed in claims 5 to 8, wherein the genes mentioned are enhanced in coryneform bacteria which already produce D-pantothenic acid.
12. The process as claimed in claim 9, wherein coryneform bacteria which already produce D-pantothenic acid and in which the pck gene is attenuated are employed.
- 25 13. An isolated polynucleotide from coryneform bacteria which lies upstream of SEQ ID No. 1 and is shown in SEQ ID No. 6.
14. An isolated polynucleotide from coryneform bacteria which lies downstream of SEQ ID No. 1 and is shown in SEQ ID No. 7.

15. An isolated polynucleotide from coryneform bacteria, containing a deletion mutation of the poxB gene shown in SEQ ID No. 12.
 16. Coryneform bacteria which carry the deletion mutation shown in SEQ ID No. 12.
- 5

Abstract

The invention relates to a process for the preparation of D-pantothenic acid by fermentation of coryneform bacteria, in which bacteria in which the nucleotide sequence which codes for pyruvate oxidase (EC 1.2.2.2) (poxB gene) is attenuated are employed, the following steps being carried out:

- a) fermentation of D-pantothenic acid-producing bacteria in which at least the gene which codes for pyruvate oxidase is attenuated.
- b) concentration of the D-pantothenic acid in the medium or in the cells of the bacteria; and
- c) isolation of the D-pantothenic acid produced.

Figure 1: Map of the plasmid pCR2.1poxBint

5

10

15

20





Figure 2:

3



15

20